

## A new method for solving conformable fractional coupled viscous Burgers' equation

Hadi Rezazadeh<sup>1</sup>, Mostafa Eslami<sup>2</sup>, Sanaz Abedin Taleghani<sup>2</sup>, Alper Korkmaz<sup>3\*</sup>

<sup>1</sup> Faculty of Engineering Technology, Amol University of Special Modern Technologies, Amol, Iran

<sup>2</sup> Department of Mathematics, Faculty of Mathematical Sciences, University of Mazandaran, Babolsar, Iran

<sup>3</sup> Department of Mathematics, ÇAKU, Çankırı, Turkey

(Received 13 Oct 2018 , accepted 06 Mar 2019)

**Abstract:** In this paper, the combined fractional Laplace transform and new Adomian decomposition method (FLTNADM) is employed for solving conformable fractional coupled Burgers equation. Because of difficulty and complex procedure of Adomian polynomials calculations, He polynomials based on homotopy perturbation method instead of Adomian polynomials, are used. The results obtained are in good agreement with the exact solution. These results show that the technique introduced here is accurate and easy to apply with less computational work than other methods.

**Keywords:** Conformable fractional Coupled Burgers equation; Fractional Laplace transform method, New Adomian decomposition method, He polynomials.

### 1 Introduction

In literature there are many definitions on fractional derivatives but the most frequently used are as below.

i) Riemann-Liouville fractional derivative is defined by [1]

$$D_t^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t (t-\varepsilon)^{-\alpha} f(\varepsilon) d\varepsilon, \quad 0 \leq \alpha < 1. \quad (1)$$

ii) Caputo fractional derivative is defined by [1]

$$D_t^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-\varepsilon)^{-\alpha} f'(\varepsilon) d\varepsilon, \quad 0 \leq \alpha < 1. \quad (2)$$

iii) Jumaris fractional derivative is defined by [2]

$$D_t^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dx} \int_0^t (t-\varepsilon)^{-\alpha} (f(\varepsilon) - f(0)) d\varepsilon, \quad 0 < \alpha < 1. \quad (3)$$

iv) He's fractional derivative is defined by [3]

$$D_t^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t (t-\varepsilon)^{-\alpha} (f_0(\varepsilon) - f(\varepsilon)) d\varepsilon, \quad 0 < \alpha < 1. \quad (4)$$

where  $f_0(\varepsilon)$  is the solution of its continuous partner of the problem with the same initial condition of the fractal partner.

$$D_t^{\alpha,\beta} f(t) = I_t^{\beta(1-\alpha)} \frac{d}{dt} I_t^{(1-\beta)(1-\alpha)} f(t), \quad t > 0, \quad 0 < \alpha < 1, \quad 0 \leq \beta \leq 1, \quad (5)$$

\*Corresponding author. E-mail address: korkmazalper@yandex.com.tr (Alper Korkmaz)

Most of the above fractional operators are defined via the fractional integrals with singular kernels which are due to their nonlocal structures. In addition, most of the nonlocal fractional derivatives don't obey the basic chain, quotient and product rules. Recently, to overcome these and other difficulties, Khalil et al. [5] introduced a new well-behaved definition of local fractional (non-integer order) derivative, called the conformable fractional derivative. The conformable fractional derivative is theoretically very easier to handle. Recently a different study appeared in literature on conformable fractional derivative. Dazhi Zhao and Maokang Luo generalized the conformable fractional derivative and give the physical interpretation of generalized conformable derivative [6]. The conformable calculus is very fascinating and is gaining an interest-see [7]-[16] and reference therein.

Mathematical models of basic flow equations describing unsteady transport problems consist of system of nonlinear parabolic and hyperbolic partial differential equations. The coupled Burgers' equations form an important class of such partial differential equations. These equations occur in a large number of physical problems such as the phenomena of turbulence, flow through a shock wave traveling in a viscous fluid, sedimentation of two inds of particles in fluid suspensions under the effect of gravity [17, 18]. Various methods have been introduced for numerical and solution of the coupled Burgers' equations. An application of mesh free interpolation method for the numerical solution of the coupled nonlinear partial differential equation is proposed in [19]. Khater et al. [20] have obtained approximate solution of the viscous coupled Burgers' equation using cubic-spline collocation method. The equation has been solved by Deghan et al. [21] using a Pade technique and Rashid et al. [22] have used Fourier Pseudo spectral method to find numerical solution of the equation. Variational iteration method has been presented for solving the coupled viscous Burgers' equation by Adbou and Soliman [23]. The exact solution of the equation has been obtained by Kaya [24] using Adomian Decomposition method and Soliman [25] presented modified extended tanh-function method to obtain its exact solution. Biazar and Ghazvini [26] proposed the homotopy perturbation method to obtain the exact solution of nonlinear Burgers' equation. Aminikhah [27] proposed a new method based on combination of the Laplace transform and new homotopy perturbation method (NHPM) first time and used it to obtain closed form solutions of coupled viscous Burgers' equation. This method has been used for solving some equations successfully [28, 29]. The main disadvantage of the Adomian method is the complex and difficult procedure for calculation the Adomian polynomials. Ghorbani [30], introduced He polynomials based on homotopy perturbation method to calculate Adomian polynomials, making the solution procedure in Adomian method remarkable simple and straightforward.

In this paper, we use the fractional Laplace transform and a new Adomian decomposition method (FLTNADM) to obtain approximate solutions with high accuracy in for homogeneous and inhomogeneous conformable fractional coupled Burgers' equations.

Rest of the paper is organized as follows. In Section 2, we will describe the conformable fractional derivative. In Section 3, the flow analysis and mathematical formulation are presented. Section 4, includes analysis of the new technique for conformable fractional coupled Burgers' equations. In Section 5, we apply the proposed method to different types of problems related to the conformable fractional coupled Burgers' equation and finally in Section 6, we summarize the results.

## 2 Conformable fractional derivative

Conformable fractional derivative of order  $\alpha$  is defined by the following definition.

**Definition 1** Let  $f : (0, \infty) \rightarrow \mathbb{R}$ , then, the conformable fractional derivative of  $f$  of order  $\alpha$  is defined as [19]

$$D_t^\alpha f(t) = \lim_{\varepsilon \rightarrow 0} \frac{f(t + \varepsilon t^{1-\alpha}) - f(t)}{\varepsilon}, \quad (6)$$

for all  $t > 0$ ,  $\alpha \in (0, 1)$ .

If  $f$  is  $\alpha$ -differentiable in some  $(0, a)$ ,  $a > 0$ , and  $\lim_{t \rightarrow 0^+} D_t^\alpha f(t)$  exists, then by definition

$$D_t^\alpha f(0) = \lim_{t \rightarrow 0^+} D_t^\alpha f(t). \quad (7)$$

This new definition of the derivative has been used to solve various problems arising in different fields of science and engineering [31–33]. The new definition satisfies the properties which given in the following theorem.

**Theorem 1** •  $D_t^\alpha (af(t) + bg(t)) = a D_t^\alpha f(t) + b D_t^\alpha g(t)$ , for all  $a, b \in \mathbb{R}$ .

- $D_t^\alpha(t^\mu) = \mu t^{\mu-\alpha}$ , for all  $\mu \in \mathbb{R}$ .
- $D_t^\alpha f(t)g(t) = f(t) D_t^\alpha g(t) + g(t) D_t^\alpha f(t)$ .
- $D_t^\alpha \frac{f(t)}{g(t)} = \frac{g(t) D_t^\alpha f(t) - f(t) D_t^\alpha g(t)}{g(t)^2}$ .

In [21] T. Abdeljawad established the chain rule for conformable fractional derivatives as following theorem.

**Theorem 2** Let  $f : (0, \infty) \rightarrow \mathbb{R}$ , be a function such that  $f$  is differentiable and also  $\alpha$ -differentiable. Let  $g$  be a function defined in the range of  $f$  and also differentiable; then, one has the following rule

$$D_t^\alpha(f \circ g)(t) = D_t^\alpha f(g(t)) D_t^\alpha g(t) g(t)^{\alpha-1}. \quad (8)$$

$$e^{\frac{1}{\alpha}t^\alpha} = \sum_{k=0}^{\infty} \frac{t^{k\alpha}}{\alpha^k k!}.$$

Now, we list here the fractional derivatives of certain functions [19]

- $D_t^\alpha(e^{\frac{1}{\alpha}t^\alpha}) = e^{\frac{1}{\alpha}t^\alpha}$ ,
- $D_t^\alpha(\sin \frac{1}{\alpha}t^\alpha) = \cos \frac{1}{\alpha}t^\alpha$ ,
- $D_t^\alpha(\cos \frac{1}{\alpha}t^\alpha) = -\sin \frac{1}{\alpha}t^\alpha$ ,
- $D_t^\alpha(\frac{1}{\alpha}t^\alpha) = 1$ .

On letting  $\alpha = 1$  in these derivatives, we get the corresponding ordinary derivatives.

**Definition 2**

$${}_t I_a^\alpha f(t) = \int_a^t \frac{f(x)}{x^{1-\alpha}} dx, \quad (9)$$

if the Riemann improper integral exists.

**Theorem 3 (Integration by parts)** [21] Let  $f, g : [0, b] \rightarrow \mathbb{R}$  be two functions such that  $fg$  is differentiable. Then

$$\int_0^b f(t) D_t^\alpha g(t) d\alpha(t) = f(t)g(t)|_0^b - \int_0^b g(t) D_t^\alpha f(t) d\alpha(t), \quad (10)$$

where  $d\alpha(t) = t^{\alpha-1} dt$ .

It is interesting to observe that the  $\alpha$ -fractional derivative and the  $\alpha$ -fractional integral are inverse of each other as given in [19].

**Theorem 4 (Inverse property)** Let  $\alpha \in (0, 1]$  and  $f$  be a continuous function such that  ${}_t I_0^\alpha f$  exists. Then

$$D_t^\alpha({}_t I_0^\alpha f)(t) = f(t), \quad \text{for } t \geq 0.$$

**Definition 3**

$$\mathcal{L}_\alpha\{f(t)\} = F_\alpha(s) = \int_0^\infty e^{-s \frac{t^\alpha}{\alpha}} f(t) d\mu(t), \quad (11)$$

where  $d\alpha(t) = t^{\alpha-1} dt$ .

Fractional Laplace transform for certain functions are presented as follows [21]

- $\mathcal{L}_\alpha\{1\} = \frac{1}{s}, \quad s > 0.$
- $\mathcal{L}_\alpha\{\frac{t^\alpha}{\alpha}\} = \frac{1}{s^2}, \quad s > 0.$
- $\mathcal{L}_\alpha\{e^{\frac{t^\alpha}{\alpha}}\} = \frac{1}{s-1}, \quad s > 1.$
- $\mathcal{L}_\alpha\{\sin \frac{1}{\alpha}t^\alpha\} = \frac{1}{s^2+1}, \quad s > 1.$
- $\mathcal{L}_\alpha\{\cos \frac{1}{\alpha}t^\alpha\} = \frac{s}{s^2+1}, \quad s > 1.$

Furthermore, using the properties of the fractional exponential function and integration by parts, we have

$$\mathcal{L}_\alpha\{D_t^\alpha(f)(t)\} = s F_\alpha(s) - f(0). \quad (12)$$

### 3 Basic Idea of the Method

In this section, we illustrate basic idea of the proposed approach. Consider the following nonlinear conformable fractional equation

$$D_x^\alpha u + N(u) = f(x), \tag{13}$$

$$D_x^\alpha U - \sum_{n=0}^{\infty} a_n \mu_n(x) + p \sum_{n=0}^{\infty} a_n \mu_n(x) + N(U) = f(x), \tag{14}$$

can be decomposed into an infinite series of polynomials given by

$$N(U) = \sum_{n=0}^{\infty} H_n(U_0, U_1, \dots, U_n), \tag{15}$$

where  $H_n(u_0, u_1, \dots, u_n)$  are called He polynomials and are defined by

$$H_n(U_0, U_1, \dots, U_n) = \left(\frac{1}{n!}\right) \left[ \frac{d^n}{dp^n} N\left(\sum_{k=0}^n p^k U_k\right) \right]_{p=0}, \quad n = 0, 1, 2, \dots \tag{16}$$

Alternatively, approximation solution of (14) can be expressed as follows

$$D_x^\alpha U - \sum_{n=0}^{\infty} a_n \mu_n(x) + p \sum_{n=0}^{\infty} a_n \mu_n(x) + \sum_{n=0}^{\infty} H_n(U_0, U_1, \dots, U_n) = f(x). \tag{17}$$

By applying fractional Laplace transform on both sides of (17), we have

$$\mathcal{L}_\alpha \left\{ D_x^\alpha U - \sum_{n=0}^{\infty} a_n \mu_n(x) + p \sum_{n=0}^{\infty} a_n \mu_n(x) + \sum_{n=0}^{\infty} H_n(U_0, U_1, \dots, U_n) - f(x) \right\} = 0, \tag{18}$$

using the differential property of fractional Laplace transform we have

$$s \mathcal{L}_\alpha \{U\} - U(0) = \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n \mu_n(x) - p \sum_{n=0}^{\infty} a_n \mu_n(x) - \sum_{n=0}^{\infty} H_n(U_0, U_1, \dots, U_n) + f(x) \right\}, \tag{19}$$

or

$$\mathcal{L}_\alpha \{U\} = \frac{1}{s} \left\{ U(0) + \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n \mu_n(x) - p \sum_{n=0}^{\infty} a_n \mu_n(x) - \sum_{n=0}^{\infty} H_n(U_0, U_1, \dots, U_n) + f(x) \right\} \right\}. \tag{20}$$

By applying inverse fractional Laplace transform on both sides of (20), we have

$$\mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} \left\{ U(0) + \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n \mu_n(x) - p \sum_{n=0}^{\infty} a_n \mu_n(x) - \sum_{n=0}^{\infty} H_n(U_0, U_1, \dots, U_n) + f(x) \right\} \right\} \right\}, \tag{21}$$

according to (21), we define

$$\begin{aligned} U_0(x) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} U(0) + \frac{1}{s} \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n \mu_n(x) \right\} \right\}, \\ U_1(x) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} \mathcal{L}_\alpha \left\{ -p \sum_{n=0}^{\infty} a_n \mu_n(x) - H_0(U_0) + f(x) \right\} \right\}, \\ &\vdots \\ U_j(x) &= \mathcal{L}_\alpha^{-1} \left\{ -\frac{1}{s} \mathcal{L}_\alpha \left\{ H_{j-1}(U_0, U_1, \dots, U_{j-1}) \right\} \right\}, \quad j = 2, 3, \dots \\ U_0(x) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} U(0) + \frac{1}{s} \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n \mu_n(x) \right\} \right\}. \end{aligned} \tag{22}$$

In this method, only the first He polynomial is calculated, and the method does not need to solve the functional equation in each iteration.

## 4 Analysis of the Method for conformable fractional coupled Burgers' equations

According to the proposed method, we consider the following conformable fractional coupled Burgers' equations

$$\begin{aligned} D_t^\alpha u - u_{xx} + \eta uu_x + \kappa(uv)_x &= f(x, t), \\ D_t^\alpha v - v_{xx} + \eta vv_x + \gamma(uv)_x &= g(x, t), \end{aligned} \tag{23}$$

$$u(x, 0) = \phi_1(x), \quad v(x, 0) = \phi_2(x), \tag{24}$$

For solving this conformable fractional equation by new method, we consider the following equation

$$\begin{aligned} D_t^\alpha U - \sum_{n=0}^{\infty} a_n(x)\mu_n(t) + p \sum_{n=0}^{\infty} a_n(x)\mu_n(t) - U_{xx} + \eta UU_x + \kappa(UV)_x &= f(x, t), \\ D_t^\alpha V - \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) + p \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) - V_{xx} + \eta VV_x + \gamma(UV)_x &= g(x, t), \end{aligned} \tag{25}$$

It is assumed that the unknown functions can be expressed by an infinite series in the form

$$\begin{aligned} U &= U_0 + U_1 + U_2 + \dots, \\ V &= V_0 + V_1 + V_2 + \dots. \end{aligned}$$

By applying fractional Laplace transform on both sides of (25), we have

$$\begin{aligned} \mathcal{L}_\alpha \left\{ D_t^\alpha U - \sum_{n=0}^{\infty} a_n(x)\mu_n(t) + p \sum_{n=0}^{\infty} a_n(x)\mu_n(t) - U_{xx} + \eta UU_x + \kappa(UV)_x - f(x, t) \right\} &= 0, \\ \mathcal{L}_\alpha \left\{ D_t^\alpha V - \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) + p \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) - V_{xx} + \eta VV_x + \gamma(UV)_x - g(x, t) \right\} &= 0, \end{aligned} \tag{26}$$

using the differential property of fractional Laplace transform we have

$$\begin{aligned} s\mathcal{L}_\alpha\{U\} - U(x, 0) &= \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n(x)\mu_n(t) - p \sum_{n=0}^{\infty} a_n(x)\mu_n(t) + U_{xx} - \eta UU_x - \kappa(UV)_x + f(x, t) \right\}, \\ s\mathcal{L}_\alpha\{V\} - V(x, 0) &= \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) - p \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) + V_{xx} - \eta VV_x - \gamma(UV)_x + g(x, t) \right\}, \end{aligned} \tag{27}$$

or

$$\begin{aligned} \mathcal{L}_\alpha\{U\} &= \frac{1}{s} \left\{ U(x, 0) + \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n(x)\mu_n(t) - p \sum_{n=0}^{\infty} a_n(x)\mu_n(t) + U_{xx} - \eta UU_x - \kappa(UV)_x + f(x, t) \right\} \right\}, \\ \mathcal{L}_\alpha\{V\} &= \frac{1}{s} \left\{ V(x, 0) + \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) - p \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) + V_{xx} - \eta VV_x - \gamma(UV)_x + g(x, t) \right\} \right\}. \end{aligned} \tag{28}$$

By applying inverse fractional Laplace transform on both sides of (28), we have

$$\begin{aligned} U(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} \left\{ U(x, 0) + \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n(x)\mu_n(t) - p \sum_{n=0}^{\infty} a_n(x)\mu_n(t) + U_{xx} - \eta \left( \sum_{n=0}^{\infty} A_n \right) - \kappa \left( \sum_{n=0}^{\infty} C_n \right) + f(x, t) \right\} \right\} \right\}, \\ V(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} \left\{ V(x, 0) + \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) - p \sum_{n=0}^{\infty} b_n(x)\lambda_n(t) + V_{xx} - \eta \left( \sum_{n=0}^{\infty} B_n \right) - \gamma \left( \sum_{n=0}^{\infty} C_n \right) + g(x, t) \right\} \right\} \right\}, \end{aligned} \tag{29}$$

$$A_n(U_0, U_1, \dots, U_n) = \left( \frac{1}{n!} \right) \left[ \frac{d^n}{d\lambda^n} \left[ N_1 \left( \sum_{k=0}^{\infty} \lambda^k U_k \right) \right] \right]_{\lambda=0},$$

$$B_n(V_0, V_1, \dots, V_n) = \left(\frac{1}{n!}\right) \left[ \frac{d^n}{d\lambda^n} [N_2 \left( \sum_{k=0}^{\infty} \lambda^k V_k \right)] \right]_{\lambda=0},$$

$$C_n(U_0, U_1, \dots, U_n, V_0, V_1, \dots, V_n) = \left(\frac{1}{n!}\right) \left[ \frac{d^n}{d\lambda^n} [N_3 \left( \sum_{k=0}^{\infty} \lambda^k U_k \cdot \sum_{k=0}^{\infty} \lambda^k V_k \right)] \right]_{\lambda=0},$$

according to (29), we define

$$\begin{cases} U_0(x, t) = \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} U(x, 0) + \frac{1}{s} \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n \mu_n(t) \right\} \right\}, \\ V_0(x, t) = \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} V(x, 0) + \frac{1}{s} \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} b_n \lambda_n(t) \right\} \right\}, \\ \left\{ \begin{aligned} U_1(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\mathcal{L}_\alpha \{-p \sum_{n=0}^{\infty} a_n \mu_n(t) + (U_0)_{xx} - \eta A_0 - \kappa C_0 + f(x, t)\}) \right\}, \\ V_1(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\mathcal{L}_\alpha \{-p \sum_{n=0}^{\infty} b_n \lambda_n(t) + (V_0)_{xx} - \eta B_0 - \gamma C_0 + g(x, t)\}) \right\}, \\ &\vdots \\ U_j(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\mathcal{L}_\alpha \{-(U_{j-1})_{xx} - \eta A_{j-1} - \kappa C_{j-1}\}) \right\}, \\ V_j(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\mathcal{L}_\alpha \{-(V_{j-1})_{xx} - \eta B_{j-1} - \gamma C_{j-1}\}) \right\}, \end{aligned} \right\} \quad j = 2, 3, \dots \end{cases} \quad (30)$$

Now, let us determine  $a_0, a_1, a_2, \dots$  so that  $U_1 = 0$ , then from (30) we have  $U_2 = U_3 = \dots = 0$ . Setting  $p = 1$ , results in the solution of equation (23) with the initial conditions as the following

$$\begin{cases} U_0(x, t) = \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} \phi_1(x) + \frac{1}{s} \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n \mu_n(t) \right\} \right\}, \\ V_0(x, t) = \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} \phi_2(x) + \frac{1}{s} \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} b_n \lambda_n(t) \right\} \right\}. \end{cases}$$

## 5 Examples

**Example 1.** Consider the following homogeneous form of a conformable fractional coupled Burgers equation

$$\begin{cases} D_t^\alpha u - u_{xx} - 2uu_x + (uv)_x = 0, \\ D_t^\alpha v - v_{xx} - 2vv_x + (uv)_x = 0, \end{cases} \quad (31)$$

subject to the initial condition

The exact solution of the equation is as  $u(x, t) = v(x, t) = e^{-\frac{1}{\alpha} t^\alpha} \sin x$ .

By using FLTNADM and setting  $U(x, 0) = V(x, 0) = \sin x$ , we have

$$\begin{cases} U_0(x, t) = \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\sin(x) + \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} a_n(x) \mu_n(t) \right\}) \right\} \\ V_0(x, t) = \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\sin(x) + \mathcal{L}_\alpha \left\{ \sum_{n=0}^{\infty} b_n(x) \lambda_n(t) \right\}) \right\} \\ \left\{ \begin{aligned} U_1(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\mathcal{L}_\alpha \{-p \sum_{n=0}^{\infty} a_n(x) \mu_n(t) + (U_0)_{xx} - 2A_0 - C_0\}) \right\}, \\ V_1(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\mathcal{L}_\alpha \{-p \sum_{n=0}^{\infty} b_n(x) \lambda_n(t) + (V_0)_{xx} - 2B_0 - C_0\}) \right\}, \\ a_0(x) &= a_2(x) = \dots = -\sin x, & a_1(x) &= a_3(x) = \dots = \sin x, \\ b_0(x) &= b_2(x) = \dots = -\sin x, & b_1(x) &= b_3(x) = \dots = \sin x, \end{aligned} \right\} \end{cases}$$




Figure 1: The surface shows the solution  $u(x, y)$  and  $v(x, y)$  for equation (31) when  $-10 \leq x \leq 10$  and  $0 < t \leq 2$ : (a)  $\alpha = 1$ , (b)  $\alpha = 0.9$

Therefore we gain the solution of equation (31) as

$$u(x, t) = U_0(x, t) = \left(1 - \frac{t^\alpha}{\alpha} + \frac{1}{2} \frac{t^{2\alpha}}{\alpha^2} - \frac{1}{6} \frac{t^{3\alpha}}{\alpha^3} + \dots\right) \sin x = e^{-\frac{1}{\alpha} t^\alpha} \sin x,$$

$$v(x, t) = V_0(x, t) = \left(1 - \frac{t^\alpha}{\alpha} + \frac{1}{2} \frac{t^{2\alpha}}{\alpha^2} - \frac{1}{6} \frac{t^{3\alpha}}{\alpha^3} + \dots\right) \sin x = e^{-\frac{1}{\alpha} t^\alpha} \sin x,$$

which are exact solution of equation (31).

**Example 2.** Consider the following non-homogeneous form of a conformable fractional coupled Burgers equation

$$\begin{aligned} D_t^\alpha u - u_{xx} + uv_x + (uv)_x &= x^2 - 2\frac{t^\alpha}{\alpha} + (2x^3 + 1)\frac{t^{2\alpha}}{2! \alpha^2}, \\ D_t^\alpha v - v_{xx} + vv_x + (uv)_x &= \frac{1}{x} - \frac{2}{x^3} \frac{t^\alpha}{\alpha} - \left(\frac{1}{x^3} - 1\right) \frac{t^{2\alpha}}{2! \alpha^2}, \end{aligned} \quad (32)$$

subject to the initial condition

The exact solution of the equation is as  $u(x, t) = x^2 \frac{t^\alpha}{\alpha}$  and  $v(x, t) = \frac{1}{x} \frac{t^\alpha}{\alpha}$ .

By using FLTNADM and setting  $U(x, 0) = V(x, 0) = 0$ , we have

$$\begin{cases} U_0(x, t) = \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (u_0 + \mathcal{L}_\alpha \{U(x, 0)\}) \right\}, \\ V_0(x, t) = \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (v_0 + \mathcal{L}_\alpha \{V(x, 0)\}) \right\}, \\ \left\{ \begin{aligned} U_1(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\mathcal{L}_\alpha \{ -p \sum_{n=0}^{\infty} a_n(x) \mu_n(t) + (U_0)_{xx} - A_0 - C_0 \right. \right. \\ &\quad \left. \left. + x^2 - 2\frac{t^\alpha}{\alpha} + (2x^3 + 1)\frac{t^{2\alpha}}{2! \alpha^2} \right\} \right\}, \\ V_1(x, t) &= \mathcal{L}_\alpha^{-1} \left\{ \frac{1}{s} (\mathcal{L}_\alpha \{ -p \sum_{n=0}^{\infty} b_n(x) \lambda_n(t) + (V_0)_{xx} - B_0 - C_0 \right. \right. \\ &\quad \left. \left. + \frac{1}{x} - \frac{2}{x^3} \frac{t^\alpha}{\alpha} - \left(\frac{1}{x^3} - 1\right) \frac{t^{2\alpha}}{2! \alpha^2} \right\} \right\}, \end{aligned} \right. \\ a_0(x) &= x^2, \quad a_1(x) = a_2(x) = a_3(x) = \dots = 0, \\ b_0(x) &= \frac{1}{x}, \quad b_1(x) = b_2(x) = b_3(x) = \dots = 0. \end{cases}$$




Figure 2: The surface shows the solution  $u(x, y)$  for equation (32) when  $-10 \leq x \leq 10$  and  $0 < t \leq 2$ : (a)  $\alpha = 1$ , (b)  $\alpha = 0.9$

Therefore we gain the solution of equation (32) as

$$u(x, t) = U_0(x, t) + U_1(x, t) + \dots = x^2 \frac{t^\alpha}{\alpha},$$

$$v(x, t) = V_0(x, t) + V_1(x, t) + \dots = \frac{1}{x} \frac{t^\alpha}{\alpha},$$

which are exact solution of equation (32).

## 6 Conclusions

In this paper, the FLTNADM, a combination of fractional Laplace transform method and new Adomian decomposition method, was applied successfully to find the analytical solution of conformable fractional coupled Burgers' equations which is a conformable fractional partial differential equation. The proposed method does not require complex computations of Adomian polynomials and only use the first term of it, so the reliability of the method and reduction in the size of computations give this method a wider applicability. The obtained results show that these approaches can solve the problem effectively and can be applied to many conformable fractional nonlinear differential equations.






Figure 3: The surface shows the solution  $v(x, y)$  for equation (32) when  $-10 \leq x \leq 10$  and  $0 < t \leq 2$ : (a)  $\alpha = 1$ , (b)  $\alpha = 0.9$ .

## References

- [1] Podlubny, I., Fractional differential equations. Vol. 198. Academic press, 1998.
- [2] Jumarie, G. (2006). Modified Riemann-Liouville derivative and fractional Taylor series of non differentiable functions further results. *Computers & Mathematics with Applications*, 51(9-10), 1367-1376.
- [3] He, J. (2014) A tutorial review on fractal spacetime and fractional calculus, *Int. J. Theor. Phys.* 53, 3698-3718.
- [4] Hilfer, R., ed. *Applications of fractional calculus in physics*. World Scientific, 2000.
- [5] Khalil, R., Al Horani, M., Yousef, A., Sababheh, M. (2014). A new definition of fractional derivative. *Journal of Computational and Applied Mathematics*, 264, 65-70.
- [6] Zhao, D., Maokang, L. (2017.) General conformable fractional derivative and its physical interpretation, *Calcolo*. 54, 903-917.
- [7] Abdeljawad, T. (2015). On conformable fractional calculus, *J. Comput. Appl. Math.* 279, 57-66.
- [8] Rezazadeh, H., Aminikhah, H., Refahi Sheikhani, A. H. (2017). Stability analysis of conformable fractional systems. *Iranian Journal of Numerical Analysis and Optimization*, 7(1), 13-32.
- [9] Aminikhah, H., Sheikhani, A. R., Rezazadeh, H. (2016). Sub-equation method for the fractional regularized long-wave equations with conformable fractional derivatives. *Scientia Iranica. Transaction B, Mechanical Engineering*, 23(3), 1048.
- [10] Rezazadeh, H., Ziabarya, B. P. (2016). Sub-equation method for the conformable fractional generalized kuramoto sivashinsky equation. *Comput. Res. Progress Appl. Sci. Eng*, 2(3), 106-109.
- [11] Eslami, M., Rezazadeh, H. (2016). The first integral method for Wu-Zhang system with conformable time-fractional derivative. *Calcolo*, 53(3), 475-485.
- [12] Hosseini, K., Mayeli, P., Ansari, R. (2017). Modified Kudryashov method for solving the conformable time-fractional Klein-Gordon equations with quadratic and cubic nonlinearities. *Optik-International Journal for Light and Electron Optics*, 130, 737-742.
- [13] Korkmaz, A. (2017). Exact solutions to  $(3+ 1)$  conformable time fractional Jimbo-Miwa, Zakharov-Kuznetsov and modified Zakharov-Kuznetsov equations. *Communications in Theoretical Physics*, 67(5), 479.
- [14] Eslami, M. (2016). Exact traveling wave solutions to the fractional coupled nonlinear Schrodinger equations. *Applied Mathematics and Computation*, 285, 141-148.
- [15] Eslami, M., Rezazadeh, H., Rezazadeh, M., Mosavi, S. S. (2017). Exact solutions to the space-time fractional Schrödinger-Hirota equation and the space-time modified KDV-Zakharov-Kuznetsov equation. *Optical and Quantum Electronics*, 49(8), 279.
- [16] Korkmaz A. (2019), Explicit Exact Solutions to Some One Dimensional Conformable Time Fractional Equations,

Waves in Random and Complex Media, 29(1), 124-137.

- [17] Burgers, J. M. (1948). A mathematical model illustrating the theory of turbulence. In *Advances in applied mechanics* (Vol. 1, pp. 171-199). Elsevier.
- [18] Cole, J. D. (1951). On a quasi-linear parabolic equation occurring in aerodynamics. *Quarterly of applied mathematics*, 9(3), 225-236.
- [19] Haq, S., Uddin, M. (2009). A meshfree interpolation method for the numerical solution of the coupled nonlinear partial differential equations. *Engineering Analysis with Boundary Elements*, 33(3), 399-409.
- [20] Khater, A. H., Temsah, R. S., Hassan, M. M. (2008). A Chebyshev spectral collocation method for solving Burgers'-type equations. *Journal of Computational and Applied Mathematics*, 222(2), 333-350.
- [21] Dehghan, M., Hamidi, A., Shakourifar, M. (2007). The solution of coupled Burgers' equations using Adomian-Pade technique. *Applied Mathematics and Computation*, 189(2), 1034-1047.
- [22] Rashid, A., Ismail, A. I. B. M. (2009). A Fourier pseudospectral method for solving coupled viscous Burgers equations. *Computational Methods in Applied Mathematics Comput. Methods Appl. Math.*, 9(4), 412-420.
- [23] Abdou, M. A., Soliman, A. A. (2005). Variational iteration method for solving Burger's and coupled Burger's equations. *Journal of Computational and Applied Mathematics*, 181(2), 245-251.
- [24] Kaya, D. (2001). An explicit solution of coupled viscous Burgers' equation by the decomposition method. *International Journal of Mathematics and Mathematical Sciences*, 27(11), 675-680.
- [25] Soliman, A. A. (2006). The modified extended tanh-function method for solving Burgers-type equations. *Physica A: Statistical Mechanics and its Applications*, 361(2), 394-404.
- [26] Biazar, J., Ghazvini, H. (2009). Exact solutions for nonlinear burgers' equation by homotopy perturbation method. *Numerical Methods for Partial Differential Equations*, 25(4), 833-842.
- [27] Aminikhah, H. (2013). An analytical approximation for coupled viscous Burgers' equation. *Applied Mathematical Modelling*, 37(8), 5979-5983.
- [28] Aminikhah, H., Sheikhan, A. R., Rezazadeh, H. (2014). An efficient method for time-fractional coupled Schrödinger system. *International Journal of Partial Differential Equations*, 2014.
- [29] Aminikhah, H., Sheikhan, A. H. R., & Rezazadeh, H. (2016). Approximate analytical solutions of distributed order fractional Riccati differential equation. *Ain Shams Engineering Journal*.
- [30] Ghorbani, A. (2009). Beyond Adomian polynomials: he polynomials. *Chaos, Solitons & Fractals*, 39(3), 1486-1492.
- [31] Osman M.S., Korkmaz A., Mirzazadeh M., Eslami M., Rezazadeh H., Zhou Q. (2018), The Unified Method for Conformable Time Fractional Schrödinger Equation with Perturbation Terms, *Chinese Journal of Physics*, 56(5), 2500-2506.
- [32] Korkmaz A., Hosseini K. (2017). Exact solutions of a nonlinear conformable time fractional parabolic equation with exponential nonlinearity using reliable methods. *Optical and Quantum Electronics*, 49(278), 1-10.
- [33] Korkmaz A. (2018). On the wave solutions of conformable fractional evolution equations. *Communications Series A1: Mathematics and Statistics*, 67(1), 68-79.
- [34] Neei J., Duan J. D., Limit set of trajectories of the coupled viscous Burgers' equations, *Appl. Math. Lett*, 11 (1998) 57-61.